WEST GEORGIA GASTROENTEROLOGY COLYTE PREP

Colon preps may leave you dehydrated, it is important to drink as much liquid as you can before, during and after the prep. Failure to take the prep as indicated may result in a poorly cleansed colon.	
NO solid foods, NO milk	
Water	
BOUILLION CUBES	
SPRITE	
7-UP	
FRUIT POPSICLES (NO RED)	
Hard Candy (NO RED)	
KOOLAID (NO RED)	

Day Prior to your Colonoscopy:

- **Step 1:** Follow the instructions to mix your prescription of Colyte.
- **Step 2: 4:00pm** Drink a glass of Colyte every 20 minutes until you finish the bottle.
- **Step 3:** Do not eat or drink anything after midnight. You may gargle but do not swallow any liquid. This includes gum, hard candy, and tobacco products.

IMPORTANT:

- 1. If you take blood pressure medicine, heart tablet or seizure medicine, you may take with a sip of water.
- 2. If you take Insulin, Take ½ of your regular dose the evening before. Do not take your diabetic pills or Insulin the morning of the test.
- 3. Your driver must stay with you the entire visit, or your procedure will be rescheduled. **After the procedure you will not be able to drive or work the remainder of the day**. Please expect your appointment to take 2 to 4 hours.

*IF YOU TAKE ASPIRIN, STOP ASPIRIN 1 DAY BEFORE PROCEDURE. IF YOU TAKE COUMADIN OR PLAVIX, YOU WILL NEED TO STOP COUMADIN OR PLAVIX 5 DAYS BEFORE PROCEDURE. IF YOU ARE A DIABETIC, DO NOT TAKE YOUR MEDICINE THE DAY OF THE PROCEDURE.